

地域包括ケア教育センター

論文

A 欧文

A-a

1. Kuba S, Ishida M, Oikawa M, Nakamura Y, Yamanouchi K, Tokunaga E, Taguchi K, Esaki T, Eguchi S, Ohno S. Aromatase inhibitors with or without luteinizing hormone-releasing hormone agonist for metastatic male breast cancer: report of four cases and review of the literature. *Breast Cancer*. 23(6):945-949, 2016. (IF: 1.412) *
2. Kuba S, Yamanouchi K, Hayashida N, Maeda S, Adachi T, Sakimura C, Kawakami F, Yano H, Matsumoto M, Otsubo R, Sato S, Fujioka H, Kuroki T, Nagayasu T, Eguchi S. Total Thyroidectomy versus Thyroid Lobectomy for Papillary Thyroid Cancer: Comparative Analysis after Propensity Score Matching: A Multicenter Study. *Int J Surg*. 2016 (IF: 2.202) *
3. Sakimura C, Eguchi S, Yamanouchi K, Minami S, Kuba S, Hayashida N, Soyama A, Hidaka M, Takatsuki M, Maeda S, Kuroki T. The first report of the thyroid function of haemophilic patients with HIV/HCV co-infection in Japan. *Haemophilia*. 22(3):e130-3. 2016 (IF: 2.498) *
4. Fukui S, Iwamoto N, Shimizu T, Umeda M, Nishino A, Koga T, Kawashiri SY, Ichinose K, Hirai Y, Tamai M, Nakamura H, Aramaki T, Iwanaga N, Izumi Y, Origuchi T, Migita K, Ueki Y, Sato S, Kawakami A. Fewer subsequent relapses and lower levels of IL-17 in Takayasu arteritis developed after the age of 40 years. *Arthritis Res Ther*. Dec 13;18(1):293. 2016 (IF: 4.707) *
5. Umeda M, Kawashiri SY, Nishino A, Koga T, Ichinose K, Michitsuji T, Shimizu T, Fukui S, Nakashima Y, Hirai Y, Iwamoto N, Aramaki T, Tamai M, Nakamura H, Origuchi T, Ueki Y, Kawakami A. Synovitis of sternoclavicular and peripheral joints can be detected by ultrasound in patients with SAPHO syndrome. *Mod Rheumatol*. Dec 9:1-5. 2016 (IF: 1.924) *
6. Kawashiri SY, Nishino A, Shimizu T, Umeda M, Fukui S, Nakashima Y, Suzuki T, Koga T, Iwamoto N, Ichinose K, Tamai M, Nakamura H, Origuchi T, Aoyagi K, Kawakami A. Ultrasound disease activity of bilateral wrist and finger joints at three months reflects the clinical response at six months of patients with rheumatoid arthritis treated with biologic disease-modifying anti-rheumatic drugs. *Mod Rheumatol*. Sep 1:1-5. 2016 (IF: 1.924) *
7. Umeda M, Ikenaga J, Koga T, Michitsuji T, Shimizu T, Fukui S, Nishino A, Nakasima Y, Kawashiri SY, Iwamoto N, Ichinose K, Hirai Y, Tamai M, Nakamura H, Origuchi T, Kawakami A. Giant Cell Arteritis which Developed after the Administration of Granulocyte-colony Stimulating Factor for Cyclic Neutropenia. *Intern Med*. 55(16):2291-4. 2016 (IF: 16.031) *
8. Kawashiri SY, Nishino A, Suzuki T, Nakashima Y, Horai Y, Ueki Y, Aramaki T, Fujikawa K, Nakashima M, Okada A, Migita K, Mizokami A, Matsuoka N, Mine M, Sakito S, Iwamoto N, Ichinose K, Tamai M, Nakamura H, Origuchi T, Aoyagi K, Eguchi K, Kawakami A. Rapid improvement of Clinical Disease Activity Index (CDAI) at 3 months predicts a preferable CDAI outcome at 1 year in active rheumatoid arthritis patients treated with tocilizumab: results from an observational investigation of daily clinical practice. *Clin Exp Rheumatol*. Sep-Oct;34(5):808-812. 2016 (IF: 2.349) *
9. Origuchi T, Fukui S, Umeda M, Nishino A, Nakashima Y, Koga T, Kawashiri SY, Iwamoto N, Ichinose K, Tamai M, Nakamura H, Kawakami A. The Severity of Takayasu Arteritis Is Associated with the HLA-B52 Allele in Japanese Patients. *Tohoku J Exp Med*. 239(1):67-72. 2016 (IF: 1.454) *
10. Nakashima Y, Tamai M, Kita J, Michitsuji T, Shimizu T, Fukui S, Umeda M, Nishino A, Suzuki T, Horai Y, Okada A, Nishimura T, Koga T, Kawashiri SY, Iwamoto N, Ichinose K, Hirai Y, Arima K, Yamasaki S, Nakamura H, Origuchi T, Takao S, Uetani M, Aoyagi K, Eguchi K, Kawakami A. Magnetic Resonance Imaging Bone Edema at Enrollment Predicts Rapid Radiographic Progression in Patients with Early RA: Results from the Nagasaki University Early Arthritis Cohort. *J Rheumatol*. 43(7):1278-84. 2016 (IF: 3.316) *
11. Ushigusa T, Ichinose K, Sato S, Michitsuji T, Shimizu T, Umeda M, Fukui S, Nishino A, Nakashima Y, Koga T, Kawashiri SY, Iwamoto N, Hirai Y, Tamai M, Nakamura H, Origuchi T, Kawakami A. Soluble α -klotho is a potential biomarker associated with neuropsychiatric systemic lupus erythematosus. *Clin Immunol*. Apr;165:29-34. 2016 (IF: 3.841) *
12. Ichinose K, Arima K, Umeda M, Fukui S, Nishino A, Nakashima Y, Suzuki T, Horai Y, Koga T, Kawashiri SY, Iwamoto N, Fujikawa K, Aramaki T, Tamai M, Nakamura H, Sato S, Origuchi T, Kawakami A. Predictors of clinical outcomes in patients with neuropsychiatric systemic lupus erythematosus. *Cytokine*. Mar;79:31-7. 2016 (IF: 2.857) *
13. Iwamoto N, Fukui S, Umeda M, Nishino A, Nakashima Y, Suzuki T, Horai Y, Nonaka F, Okada A, Koga T, Kawashiri SY, Fujikawa K, Aramaki T, Ichinose K, Hirai Y, Tamai M, Nakamura H, Terada K, Nakashima M, Mizokami A, Origuchi T, Eguchi K, Ueki Y, Kawakami A. Evaluation of switching from intravenous to subcutaneous formulation of tocilizumab in patients with rheumatoid arthritis. *Mod Rheumatol*. Sep;26(5):662-6. 2016 (IF: 1.924) *
14. Ichinose K, Ushigusa T, Nishino A, Nakashima Y, Suzuki T, Horai Y, Koga T, Kawashiri SY, Iwamoto N, Tamai M, Arima K, Nakamura H, Obata Y, Yamamoto K, Origuchi T, Nishino T, Kawakami A, Tsokos GC. Lupus Nephritis IgG Induction of Calcium/Calmodulin-Dependent Protein Kinase IV Expression in Podocytes and Alteration of Their Function. *Arthritis Rheumatol*. Apr;68(4):944-52. 2016 (IF: 1.924) *

15. Kawashiri SY, Nishino A, Nakamura H, Origuchi T, Aoyagi K, Kawakami A. The stiff elastographic change of submandibular glands rapidly improves after the introduction of glucocorticoid treatment in patients with IgG4-related dacryoadenitis and sialoadenitis. *Mod Rheumatol.* 26(3):463-4. 2016 (IF: 1.924) *
16. Umeda M, Ichinose K, Okada A, Nishino A, Aramaki T, Iwanaga N, Terada K, Nakamura H, Ueki Y, Kawakami A. A rare case of hemorrhagic cystitis complicated with thrombocytopenia and hemophagocytic syndrome associated with BK virus, under immunosuppressive treatment of systemic lupus erythematosus. *Mod Rheumatol.* 26(3):467-9 2016 (IF: 1.924) *
17. Nishino A, Kawashiri SY, Shimizu T, Umeda M, Fukui S, Koga T, Iwamoto N, Ichinose K, Tamai M, Nakamura H, Origuchi T, Nagata Y, Maeda T, Aoyagi K, Kawakami A. Assessment of both articular synovitis and tenosynovitis by ultrasound is useful for evaluations of hand dysfunction in early rheumatoid arthritis patients. *Mod Rheumatol.* Nov 15:1-4. 2016 (IF: 1.924) *

A-e

1. Fukui S, Iwamoto N, Umeda M, Nishino A, Nakashima Y, Koga T, Kawashiri SY, Ichinose K, Hirai Y, Tamai M, Nakamura H, Origuchi T, Sato S, Kawakami A. ④ Antineutrophilic cytoplasmic antibody-associated vasculitis with hypocomplementemia has a higher incidence of serious organ damage and a poor prognosis. *Medicine (Baltimore).* Sep;95(37):e4871. 2016

B 邦文

B-a

1. 久芳さやか、伊東昌子、松坂雄亮、牟田久美子、濱口由子、相良郁子、石居公之、江口 晋、永田康浩：キャリアを継続するために卒前教育のできる事とは？ —医学生におけるワークライフバランス、将来設計の意識調査より— *長崎医学会雑誌* 91(1):1-6.2016.3

B-b

1. 永田康浩 複合がん免疫療法—外科的視点から *Cancer Board Square* Vol2, no3, 2016

研究業績集計表 教室等名：地域包括ケア教育センター

論文数一覧

	A-a	A-b	A-c	A-d	A-e	合計	SCI	B-a	B-b	B-c	B-d	B-e	合計	総計
2016	17	0	0	0	1	18	17	1	1	0	0	0	2	20

学会発表数一覧

	A-a	A-b		合計	B-a	B-b		合計	総計
		シンポジウム	学会			シンポジウム	学会		
2016	0	0	0	0	0	0	15	15	15

論文総数に係る教員生産係数一覧

	$\frac{\text{欧文論文総数}}{\text{論文総数}}$	教員生産係数 (欧文論文)	$\frac{\text{SCI掲載論文数}}{\text{欧文論文総数}}$	教員生産係数 (SCI掲載論文)
2016	0.9	4.5	0.944	4.25

Impact factor 値一覧

	Impact factor	教員当たり Impact factor	論文当たり Impact factor
2016	54.135	13.534	3.184